arXiv:math/9905141v1 [math.QA] 21 May 1999

Two-Dimensional Centrally Extended Quantum Galilei Groups and their Algebras

Anna Opanowicz*
Department of Theoretical Physics II
University of Łódź
Pomorska 149/153, 90-236 Łódź, Poland

Abstract

All deformations of two dimensional centrally extended Galilei group are classified. The corresponding quantum Lie algebras are found.

---

[*] Supported by Łódź University Grant No. 623

1 Introduction

In this paper we continue the study of the problem of deformations of space-time symmetry groups. In recent paper [1] we classified all nonequivalent Lie-Poisson structures for centrally extended two-dimensional Galilei group. In the present paper we quantize these structures by replacing the Poisson brackets by commutators (this method has been used in Ref.[2] to obtain the deformations of two-dimensional Galilei group and in Ref.[3] to construct quantum group dual to the deformed $e_q(2)$). As a result we obtain Hopf algebras and find by duality relations the corresponding quantum Lie algebras.

2 Deformations of two-dimensional centrally extended Galilei group

The classical centrally extended Galilei group is defined as a set of elements

$$g = (m, \tau, v, a) \tag{1}$$

where τ is time translation, a and v are space translation and Galilean boost, respectively, subject to the following multiplication law:

$$g'g = (m' + m - \frac{1}{2}v'^2\tau - av', \tau' + \tau, v' + v, a' + a + \tau v') \tag{2}$$

The multiplication rule defines coproducts of m, v, τ and a:

$$\begin{array}{l} \triangle(m) = m \otimes I + I \otimes m - \frac{1}{2}v^2 \otimes \tau - v \otimes a \\ \triangle(\tau) = \tau \otimes I + I \otimes \tau \\ \triangle(a) = a \otimes I + I \otimes a + v \otimes \tau \\ \triangle(v) = v \otimes I + I \otimes v \end{array} \tag{3}$$

The antipode and counit can be also read off from Eq.(2)

$$\begin{array}{l} S(m) = -m + \frac{1}{2}v^2\tau - av \\ S(\tau) = -\tau \\ S(a) = -a + v\tau \\ S(v) = -v, \end{array} \tag{4}$$

$$\varepsilon(m) = 0,\ \varepsilon(\tau) = 0,\ \varepsilon(a) = 0,\ \varepsilon(v) = 0 \tag{5}$$

Galilei group is a real Lie group. The reality condition can be described by introducing the following *-structure:

$$m^* = m,\ \tau^* = \tau,\ a^* = a,\ v^* = v \tag{6}$$

The starting point to obtain two-dimensional quantum Galilei group are the Lie-Poisson structures on it. In Ref.[1] we have found all relevant Poisson structures.

In order to obtain the corresponding quantum group we make a replacement

$$\{,\} = \frac{1}{i\kappa}[,] \tag{7}$$

Where κ is an arbitrary parameter. The quantization procedure applied to the Lie-Poisson structures classified in Ref.[1] yield the noncommutative structures listed in Table 1.

Table 1

	$[v, a]$	$[v, m]$	$[\tau, a]$	$[\tau, m]$	$[a, m]$	
1			$i\frac{1}{\chi}v$	$-\frac{1}{2}i\frac{1}{\chi}v^2$	$-\frac{1}{6}i\frac{1}{\chi}v^3$	$\frac{1}{\chi} = \frac{{\tau_0}^2}{\kappa}$
2		$i\frac{1}{\chi}v$			$i\frac{1}{\chi}a$	$\frac{1}{\chi} = \frac{{v_0}^2\tau_0}{\kappa}$
3				$i\frac{1}{\chi}v$	$\frac{1}{2}i\frac{1}{\chi}v^2$	$\frac{1}{\chi} = \frac{v_0{\tau_0}^2}{\kappa}$
4					$-i\frac{1}{\chi}\tau$	$\frac{1}{\chi} = \frac{{v_0}^3\tau_0}{\kappa}$
5					$i\frac{1}{\chi}v$	$\frac{1}{\chi} = \frac{{v_0}^2{\tau_0}^2}{\kappa}$
6		$i\frac{1}{\chi_1}v$			$i\frac{1}{\chi_1}a + i\frac{1}{\chi_2}v$	$\frac{1}{\chi_1} = \frac{{v_0}^2\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{{v_0}^2{\tau_0}^2}{\kappa}$
7		$i\frac{1}{\chi_1}v$	$i\frac{1}{\chi_2}v$	$-\frac{1}{2}i\frac{1}{\chi_2}v^2$	$i\frac{1}{\chi_1}a - \frac{1}{6}i\frac{1}{\chi_2}v^3$	$\frac{1}{\chi_1} = \frac{{v_0}^2\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{{\tau_0}^2}{\kappa}$
8				$i\frac{1}{\chi_1}v$	$\frac{1}{2}i\frac{1}{\chi_1}v^2 - i\frac{1}{\chi_2}\tau$	$\frac{1}{\chi_1} = \frac{v_0{\tau_0}^2}{\kappa}, \frac{1}{\chi_2} = \frac{{v_0}^3\tau_0}{\kappa}$
9	$i\frac{1}{\chi_1}v$	$-\frac{1}{2}i\frac{1}{\chi_1}v^2$		$i\frac{1}{\chi_2}v$	$i\frac{1}{\chi_3}v - i\frac{1}{\chi_1}m + \frac{1}{2}i\frac{1}{\chi_2}v^2$	$\frac{1}{\chi_1} = \frac{v_0\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{v_0{\tau_0}^2}{\kappa}, \frac{1}{\chi_3} = \frac{\varepsilon{v_0}^2{\tau_0}^2}{\kappa}$
10			$i\frac{1}{\chi_1}v$	$-\frac{1}{2}i\frac{1}{\chi_1}v^2$	$-i\frac{1}{\chi_2}\tau - \frac{1}{6}i\frac{1}{\chi_1}v^3$	$\frac{1}{\chi_1} = \frac{{\tau_0}^2}{\kappa}, \frac{1}{\chi_2} = \frac{{v_0}^3\tau_0}{\kappa}$
11				$-i\frac{1}{\chi_1}\tau$	$-i\frac{1}{\chi_2}\tau - i\frac{1}{\chi_1}a$	$\frac{1}{\chi_1} = \frac{{v_0}^2\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{{v_0}^3\tau_0}{\kappa}$
12		$i\frac{1}{\chi_1}v$		$-i\frac{1}{\chi_2}\tau$	$i(\frac{1}{\chi_1} - \frac{1}{\chi_2})a$	$\frac{1}{\chi_1} = \frac{{v_0}^2\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{\varepsilon{v_0}^2\tau_0}{\kappa}$
13		$i\frac{1}{\chi_1}v$		$i\frac{1}{\chi_1}\tau + i\frac{1}{\chi_2}v$	$2i\frac{1}{\chi_1}a + \frac{1}{2}i\frac{1}{\chi_2}v^2$	$\frac{1}{\chi_1} = \frac{{v_0}^2\tau_0}{\kappa}, \frac{1}{\chi_2} = \frac{v_0{\tau_0}^2}{\kappa}$
14		$-i\frac{1}{\chi_1}\tau$		$-i\frac{1}{\chi_2}\tau$	$-i\frac{1}{\chi_2}a - \frac{1}{2}i\frac{1}{\chi_1}\tau^2 + i\frac{1}{\chi_3}v$	$\frac{1}{\chi_1} = \frac{{v_0}^3}{\kappa}, \frac{1}{\chi_3} = \frac{{v_0}^2{\tau_0}^2}{\kappa}, \frac{1}{\chi_2} = \frac{\varepsilon{v_0}^2\tau_0}{\kappa}$
15		$-i\frac{1}{\chi_1}\tau$		$-i\frac{1}{\chi_2}\tau + i\frac{1}{\chi_3}v$	$-i\frac{1}{\chi_2}a - \frac{1}{2}i\frac{1}{\chi_1}\tau^2 + \frac{1}{2}i\frac{1}{\chi_3}v^2$	$\frac{1}{\chi_1} = \frac{{v_0}^3}{\kappa}, \frac{1}{\chi_3} = \frac{v_0{\tau_0}^2}{\kappa}, \frac{1}{\chi_2} = \frac{\varepsilon{v_0}^2\tau_0}{\kappa}$
16			$-i\frac{1}{\chi}\tau$	$i\frac{1}{\chi}a$	$-i\frac{1}{\chi}m$	$\frac{1}{\chi} = \frac{v_0\tau_0}{\kappa}$

and, for all cases:

$$[v, \tau] = 0 \tag{8}$$

We have checked that the comutators listed above satisfy the Jacobi identities. We have also verified that the equations (3), (4), (5), (6) together with Table 1 and (8) define a *-Hopf algebras which provide the deformations of two-dimensional centrally extended Galilei group.

3 The quantum Lie algebra and coalgebra

In this section we find all quantum Lie algebras corresponding to the quantum Galilei groups defined in Sec.2. To this end we use the Hopf algebra duality rules. On the classsical level the generators of Lie algebra of Galilei group can be defined by the following global parametrization of group element

$$g = e^{imM} e^{-i\tau H} e^{iaP} e^{ivK} \tag{9}$$

We adopt here this definition as well as classical duality relation

$$< \varphi, X >= -i\frac{d}{dt}\varphi(e^{itX})|_{t=o} \tag{10}$$

The group algebra is generated by the set of elements of the form

$$\varphi = m^{\alpha}\tau^{\beta}a^{\gamma}v^{\sigma} \tag{11}$$

where $\alpha, \beta, \gamma, \sigma \geq 1$. By applying the classical duality rules we obtain

$$\begin{array}{l} < M, m^{\alpha}\tau^{\beta}a^{\gamma}v^{\sigma} >= -i\delta_{1\alpha}\delta_{0\beta}\delta_{0\gamma}\delta_{0\sigma} \\ < H, m^{\alpha}\tau^{\beta}a^{\gamma}v^{\sigma} >= i\delta_{0\alpha}\delta_{1\beta}\delta_{0\gamma}\delta_{0\sigma} \\ < P, m^{\alpha}\tau^{\beta}a^{\gamma}v^{\sigma} >= -i\delta_{0\alpha}\delta_{0\beta}\delta_{1\gamma}\delta_{0\sigma} \\ < K, m^{\alpha}\tau^{\beta}a^{\gamma}v^{\sigma} >= -i\delta_{0\alpha}\delta_{0\beta}\delta_{0\gamma}\delta_{1\sigma}. \end{array} \tag{12}$$

In order to define dual algebra structures we use the duality relation:

$$\begin{array}{l} < \varphi, XY >=< \triangle\varphi, X \otimes Y > \\ < \varphi\psi, X >=< \varphi \otimes \psi, \triangle(X) > \end{array} \tag{13}$$

The *-structure on dual Hopf algebra can be defined by the formula

$$< X^*, \varphi >=< X, S^{-1}(\varphi^*) > \tag{14}$$

provided the following relation holds

$$S^{-1}(X) = [S(X^*)]^* \tag{15}$$

In order to take care about all elements of the form (11) let us introduce the generating function depending on four real parameters $\mu, \nu, \varrho, \kappa$

$$\varphi(\mu, \nu, \rho, \kappa) =< \varphi, e^{\mu m} e^{\nu \tau} e^{\rho a} e^{\kappa v} > \tag{16}$$

Then we have

$$\begin{array}{l} < M, e^{\mu m} e^{\nu \tau} e^{\rho a} e^{\kappa v} >= -i\mu \\ < H, e^{\mu m} e^{\nu \tau} e^{\rho a} e^{\kappa v} >= i\nu \\ < P, e^{\mu m} e^{\nu \tau} e^{\rho a} e^{\kappa v} >= -i\varrho \\ < K, e^{\mu m} e^{\nu \tau} e^{\rho a} e^{\kappa v} >= -i\kappa \end{array} \tag{17}$$

The duality rules (13) and long and tedious calculations lead us to the following structure of quantum Lie algebras

$$\begin{array}{l} 1. \\ \;\; [K, P] = iM \\ \;\; [K, H] = iP \\ \\ \;\; \Delta M = I \otimes M + M \otimes I \\ \;\; \Delta H = I \otimes H + H \otimes I \\ \;\; \Delta P = I \otimes P + P \otimes I \\ \;\; \Delta K = I \otimes K + K \otimes I - \frac{1}{\chi} P \otimes H \\ \\ \;\; S(M) = -M \\ \;\; S(H) = -H \\ \;\; S(P) = -P \\ \;\; S(K) = -K - \frac{1}{\chi} PH \end{array} \tag{18}$$

2.

$$\begin{aligned}
&[K,H] = iP\\
&[K,P] = \tfrac{1}{2} i\chi(1 - e^{-2\frac{1}{\chi}M})\\
&\\
&\Delta M = I \otimes M + M \otimes I\\
&\Delta H = I \otimes H + H \otimes I\\
&\Delta P = I \otimes P + P \otimes e^{-\frac{1}{\chi}M}\\
&\Delta K = I \otimes K + K \otimes e^{-\frac{1}{\chi}M}\\
&\\
&S(M) = -M\\
&S(H) = -H\\
&S(P) = -Pe^{\frac{1}{\chi}M}\\
&S(K) = -Ke^{\frac{1}{\chi}M}
\end{aligned} \tag{19}$$

3.

$$\begin{aligned}
&[K,P] = iM\\
&[K,H] = iP\\
&\\
&\Delta M = I \otimes M + M \otimes I\\
&\Delta H = I \otimes H + H \otimes I\\
&\Delta P = I \otimes P + P \otimes I\\
&\Delta K = I \otimes K + K \otimes I + \frac{1}{\chi} H \otimes M\\
&\\
&S(M) = -M\\
&S(H) = -H\\
&S(P) = -P\\
&S(K) = -K - \frac{1}{\chi} HM
\end{aligned} \tag{20}$$

4.

$$\begin{aligned}
&[K,P]=iM\\
&[K,H]=iP-\tfrac{1}{2}i\tfrac{1}{\chi}M^2\\
\\
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes I-\tfrac{1}{\chi}P\otimes M\\
&\Delta P=I\otimes P+P\otimes I\\
&\Delta K=I\otimes K+K\otimes I\\
\\
&S(M)=-M\\
&S(H)=-H-\tfrac{1}{\chi}PM\\
&S(P)=-P\\
&S(K)=-K
\end{aligned}\tag{21}$$

5.

$$\begin{aligned}
&[K,P]=iM\\
&[K,H]=iP\\
\\
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes I\\
&\Delta P=I\otimes P+P\otimes I\\
&\Delta K=I\otimes K+K\otimes I-\tfrac{1}{\chi}P\otimes M\\
\\
&S(M)=-M\\
&S(H)=-H\\
&S(P)=-P\\
&S(K)=-K-\tfrac{1}{\chi}PM
\end{aligned}\tag{22}$$

6.

$$
\begin{aligned}
&[K,H] = iP\\
&[K,P] = \tfrac{1}{2} i\chi_1 (1 - e^{-2\frac{1}{\chi_1}M})
\end{aligned}
$$

$$
\begin{aligned}
&\Delta M = I \otimes M + M \otimes I\\
&\Delta H = I \otimes H + H \otimes I\\
&\Delta P = I \otimes P + P \otimes e^{-\frac{1}{\chi_1}M}\\
&\Delta K = I \otimes K + K \otimes e^{-\frac{1}{\chi_1}M} - \tfrac{1}{\chi_2} P \otimes M e^{-\frac{1}{\chi_1}M}
\end{aligned}
\tag{23}
$$

$$
\begin{aligned}
&S(M) = -M\\
&S(H) = -H\\
&S(P) = -Pe^{\frac{1}{\chi_1}M}\\
&S(K) = -Ke^{\frac{1}{\chi_1}M} - \tfrac{1}{\chi_2} PMe^{\frac{1}{\chi_1}M}
\end{aligned}
$$

7.

$$
\begin{aligned}
&[K,H] = iP\\
&[K,P] = \tfrac{1}{2} i\chi_1 (1 - e^{-2\frac{1}{\chi_1}M})
\end{aligned}
$$

$$
\begin{aligned}
&\Delta M = I \otimes M + M \otimes I\\
&\Delta H = I \otimes H + H \otimes I\\
&\Delta P = I \otimes P + P \otimes e^{-\frac{1}{\chi_1}M}\\
&\Delta K = I \otimes K + K \otimes e^{-\frac{1}{\chi_1}M} - \tfrac{1}{\chi_2} P \otimes H e^{-\frac{1}{\chi_1}M}
\end{aligned}
\tag{24}
$$

$$
\begin{aligned}
&S(M) = -M\\
&S(H) = -H\\
&S(P) = -Pe^{\frac{1}{\chi_1}M}\\
&S(K) = -Ke^{\frac{1}{\chi_1}M} - \tfrac{1}{\chi_2} PHe^{\frac{1}{\chi_1}M}
\end{aligned}
$$

8.

$$[K,P]=iM$$
$$[K,H]=iP-\frac{1}{2}i\frac{1}{\chi_2}M^2$$

$$\begin{aligned}
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes I-\frac{1}{\chi_2}P\otimes M\\
&\Delta P=I\otimes P+P\otimes I\\
&\Delta K=I\otimes K+K\otimes I+\frac{1}{\chi_1}H\otimes M-\frac{1}{2}\frac{1}{\chi_1}\frac{1}{\chi_2}P\otimes M^2
\end{aligned}\tag{25}$$

$$S(M)=-M$$
$$S(H)=-H-\frac{1}{\chi_2}PM$$
$$S(P)=-P$$
$$S(K)=-K+\frac{1}{\chi_1}HM+\frac{1}{2}\frac{1}{\chi_1}\frac{1}{\chi_2}PM^2$$

9.

$$[K,M]=\frac{1}{2}i\frac{1}{\chi_1}M^2$$
$$[K,P]=iM$$
$$[K,H]=-i\chi_1(1-e^{\frac{1}{\chi_1}P})$$

$$\begin{aligned}
&\Delta M=e^{\frac{1}{\chi_1}P}\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes I\\
&\Delta P=I\otimes P+P\otimes I\\
&\Delta K=e^{\frac{1}{\chi_1}P}\otimes K+K\otimes I+\frac{1}{\chi_2}He^{\frac{1}{\chi_1}P}\otimes M-\frac{1}{\chi_3}Pe^{\frac{1}{\chi_1}P}\otimes M
\end{aligned}\tag{26}$$

$$S(M)=-Me^{-\frac{1}{\chi_1}P}$$
$$S(H)=-H$$
$$S(P)=-P$$
$$S(K)=-Ke^{-\frac{1}{\chi_1}P}+\frac{1}{\chi_2}HMe^{-\frac{1}{\chi_1}P}-\frac{1}{\chi_3}PMe^{-\frac{1}{\chi_1}P}$$

10.

$$\begin{aligned}
&[K,P]=iM\\
&[K,H]=iP-\tfrac{1}{2}i\tfrac{1}{\chi_2}M^2
\end{aligned}$$

$$\begin{aligned}
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes I-\tfrac{1}{\chi_2}P\otimes M\\
&\Delta P=I\otimes P+P\otimes I\\
&\Delta K=I\otimes K+K\otimes I+\tfrac{1}{2}\tfrac{1}{\chi_1}\tfrac{1}{\chi_2}P^2\otimes M-\tfrac{1}{\chi_1}P\otimes H
\end{aligned}\tag{27}$$

$$\begin{aligned}
&S(M)=-M\\
&S(H)=-H-\tfrac{1}{\chi_2}PM\\
&S(P)=-P\\
&S(K)=-K-\tfrac{1}{\chi_1}PM-\tfrac{1}{2}\tfrac{1}{\chi_1}\tfrac{1}{\chi_2}MP^2
\end{aligned}$$

11.

$$\begin{aligned}
&[K,H]=iP-i\tfrac{\chi_1}{\chi_2}Me^{\frac{1}{\chi_1}M}+i\tfrac{{\chi_1}^2}{\chi_2}(e^{\frac{1}{\chi_1}M}-1)\\
&[K,P]=-i\chi_1(e^{\frac{1}{\chi_1}M}-1)
\end{aligned}$$

$$\begin{aligned}
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes e^{\frac{1}{\chi_1}M}-\tfrac{1}{\chi_2}P\otimes Me^{\frac{1}{\chi_1}M}\\
&\Delta P=I\otimes P+P\otimes e^{\frac{1}{\chi_1}M}\\
&\Delta K=I\otimes K+K\otimes I
\end{aligned}\tag{28}$$

$$\begin{aligned}
&S(M)=-M\\
&S(H)=-He^{-\frac{1}{\chi_1}M}-\tfrac{1}{\chi_2}PMe^{-\frac{1}{\chi_1}M}\\
&S(P)=-Pe^{-\frac{1}{\chi_1}M}\\
&S(K)=-K
\end{aligned}$$

12.

$$
\begin{aligned}
&[K,H]=iP\\
&[K,P]=-\frac{i}{(2\frac{1}{\chi_1}-\frac{1}{\chi_2})}(1-e^{-(2\frac{1}{\chi_1}-\frac{1}{\chi_2})M})\\
&\\
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes e^{\frac{1}{\chi_2}M}\\
&\Delta P=I\otimes P+P\otimes e^{(\frac{1}{\chi_2}-\frac{1}{\chi_1})M}\\
&\Delta K=I\otimes K+K\otimes e^{-\frac{1}{\chi_1}M}\\
&\\
&S(M)=-M\\
&S(H)=-He^{-\frac{1}{\chi_2}M}\\
&S(P)=-Pe^{(\frac{1}{\chi_1}-\frac{1}{\chi_2})M}\\
&S(K)=-Ke^{\frac{1}{\chi_1}M}
\end{aligned}
\tag{29}
$$

13.

$$
\begin{aligned}
&[K,H]=iP\\
&[K,P]=\tfrac{1}{3}i\chi_1(1-e^{-3\frac{1}{\chi_1}M})\\
&\\
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes e^{-\frac{1}{\chi_1}M}\\
&\Delta P=I\otimes P+P\otimes e^{-2\frac{1}{\chi_1}M}\\
&\Delta K=I\otimes K+K\otimes e^{-\frac{1}{\chi_1}M}+\tfrac{1}{\chi_2}H\otimes Me^{-\frac{1}{\chi_1}M}\\
&\\
&S(M)=-M\\
&S(H)=-He^{\frac{1}{\chi_1}M}\\
&S(P)=-Pe^{2\frac{1}{\chi_1}M}\\
&S(K)=-Ke^{\frac{1}{\chi_1}M}+\tfrac{1}{\chi_2}HMe^{\frac{1}{\chi_1}M}
\end{aligned}
\tag{30}
$$

14.

$$[K,H] = iP + i\frac{{\chi_2}^2}{\chi_1\chi_3}Me^{\frac{1}{\chi_2}M} - \frac{1}{2}i\frac{{\chi_2}^3}{\chi_1\chi_3}(e^{2\frac{1}{\chi_2}M} - 1)$$

$$[K,P] = i\chi_2(e^{\frac{1}{\chi_2}M} - 1)$$

$$[H,P] = -\frac{1}{2}i\frac{{\chi_2}^2}{\chi_1}(e^{2\frac{1}{\chi_2}M} - 1) + i\frac{{\chi_2}^2}{\chi_1}(e^{\frac{1}{\chi_2}M} - 1)$$

$$\begin{aligned}
&\Delta M = I \otimes M + M \otimes I \\
&\Delta H = I \otimes H + H \otimes e^{\frac{1}{\chi_2}M} - \frac{\chi_2}{\chi_1}K \otimes e^{\frac{1}{\chi_2}M} + \frac{\chi_2}{\chi_1}K \otimes I \\
&+\frac{\chi_2}{\chi_1\chi_3}P \otimes Me^{\frac{1}{\chi_2}M} + \frac{{\chi_2}^2}{\chi_1\chi_3}P \otimes I - \frac{{\chi_2}^2}{\chi_1\chi_3}P \otimes e^{\frac{1}{\chi_2}M} \\
&\Delta P = I \otimes P + P \otimes e^{\frac{1}{\chi_2}M} \\
&\Delta K = I \otimes K + K \otimes I + \frac{\chi_2}{\chi_3}P \otimes I - \frac{\chi_2}{\chi_3}P \otimes e^{\frac{1}{\chi_2}M}
\end{aligned} \tag{31}$$

$$\begin{aligned}
&S(M) = -M \\
&S(H) = -He^{-\frac{1}{\chi_2}M} - \frac{\chi_2}{\chi_1}K + \frac{\chi_2}{\chi_1}Ke^{-\frac{1}{\chi_2}M} + \frac{{\chi_2}^2}{\chi_1\chi_3}Pe^{-\frac{1}{\chi_2}M} \\
&-\frac{{\chi_2}^2}{\chi_1\chi_3}P + \frac{\chi_2}{\chi_1\chi_3}PMe^{-\frac{1}{\chi_2}M} \\
&S(P) = -Pe^{-\frac{1}{\chi_2}M} \\
&S(K) = -K + \frac{\chi_2}{\chi_3}Pe^{-\frac{1}{\chi_2}M} - \frac{\chi_2}{\chi_3}P
\end{aligned}$$

15.

$$
\begin{aligned}
&[K,H]=iP\\
&[K,P]=\frac{i\left(-\frac{1}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}{\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\left(\frac{3}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}\left(e^{(\frac{3}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}})M}-1\right)\\
&+\frac{i\left(\frac{1}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}{\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\left(\frac{3}{2}\frac{1}{\chi_2}-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}\left(e^{(\frac{3}{2}\frac{1}{\chi_2}-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}})M}-1\right)\\
&[H,P]=\frac{i}{\chi_1\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\left(\frac{3}{2}\frac{1}{\chi_2}-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}\left(e^{(\frac{3}{2}\frac{1}{\chi_2}-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}})M}-1\right)\\
&-\frac{i}{\chi_1\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\left(\frac{3}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}\right)}\left(e^{(\frac{3}{2}\frac{1}{\chi_2}+\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}})M}-1\right)\\
\\
&\Delta M=I\otimes M+M\otimes I\\
&\Delta H=I\otimes H+H\otimes e^{\frac{1}{2}\frac{1}{\chi_2}M}[\cosh(-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}M)\\
&-\frac{1}{\chi_2\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}\sinh(-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}M)]\\
&+2\frac{1}{\chi_1\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}K\otimes e^{\frac{1}{2}\frac{1}{\chi_2}M}\sinh(-\frac{1}{2}\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}M)
\end{aligned}
\qquad (32)
$$

$$
\begin{aligned}
&\Delta P = I \otimes P + P \otimes e^{\frac{1}{\chi_2}M} \\
&\Delta K = I \otimes K + K \otimes e^{\frac{1}{2}\frac{1}{\chi_2}M}[\cosh(-\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M) \\
&+\frac{1}{\chi_2\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}\sinh(-\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M)] \\
&-2\frac{1}{\chi_3\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}H \otimes e^{\frac{1}{2}\frac{1}{\chi_2}M}\sinh(-\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M) \\
\\
&S(M) = -M \\
&S(H) = -He^{-\frac{1}{2}\frac{1}{\chi_2}M}[\cosh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M) \\
&-\frac{1}{\chi_2\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}\sinh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M)] \\
&-2\frac{1}{\chi_1\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}Ke^{-\frac{1}{2}\frac{1}{\chi_2}M}\sinh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M) \\
&S(P) = -Pe^{-\frac{1}{\chi_2}M} \\
&S(K) = -Ke^{-\frac{1}{2}\frac{1}{\chi_2}M}[\cosh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M) \\
&+\frac{1}{\chi_2\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}\sinh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M)] \\
&+2\frac{1}{\chi_3\sqrt{\frac{1}{{\chi_2}^2}-4\frac{1}{\chi_1}\frac{1}{\chi_3}}}He^{-\frac{1}{2}\frac{1}{\chi_2}M}\sinh(\tfrac{1}{2}\sqrt{\tfrac{1}{{\chi_2}^2} - 4\tfrac{1}{\chi_1}\tfrac{1}{\chi_3}}M)
\end{aligned}
\tag{33}
$$

16.

$$\begin{array}{l}
[K, M] = \frac{1}{2} i \frac{1}{\chi} M^2 \\
[K, P] = iM \\
[K, H] = i\chi(1 - e^{-\frac{1}{\chi}P}) - i\frac{1}{\chi}MH \\
\\
\Delta M = M \otimes I + \frac{e^{\frac{1}{\chi}P} \otimes M}{1 - \frac{1}{2}\frac{1}{\chi^2} H e^{\frac{1}{\chi}P} \otimes M} \\
\Delta H = e^{-\frac{1}{\chi}P} \otimes H + H \otimes I - \frac{1}{\chi^2} H \otimes MH + \frac{1}{4}\frac{1}{\chi^4} H^2 e^{\frac{1}{\chi}P} \otimes M^2 H \\
-\frac{1}{2}\frac{1}{\chi^2} H^2 e^{\frac{1}{\chi}P} \otimes M \\
\Delta P = I \otimes P + P \otimes I - 2\chi \ln(1 - \frac{1}{2}\frac{1}{\chi^2} H e^{\frac{1}{\chi}P} \otimes M) \\
\Delta K = I \otimes K + K \otimes I \\
\\
S(M) = \frac{-Me^{-\frac{1}{\chi}P}}{1 - \frac{1}{2}\frac{1}{\chi^2} MH} \\
S(H) = -He^{\frac{1}{\chi}P} + \frac{1}{2}\frac{1}{\chi^2} MH^2 e^{\frac{1}{\chi}P} \\
S(P) = -P + 2\chi \ln\left(\frac{1}{1 - \frac{1}{2}\frac{1}{\chi^2} HM}\right) \\
S(K) = -K
\end{array} \tag{34}$$

The counits for all cases take the form

$$\varepsilon(M) = 0,\ \varepsilon(H) = 0,\ \varepsilon(P) = 0,\ \varepsilon(K) = 0 \tag{35}$$

4 Conclusions

We obtained a number of, in general multiparameter, deformations of two-dimensional centrally extended Galilei group. They are described in Table 1 together with Eqs. (3), (4), (5), (6) and (8). The corresponding quantum Lie algebras have been also found. They are listed in Eqs.(18) - (35). Some of the resulting structures seem to be quite interesting and deserve more detailed study which will be the subject of subsequent publications.

5 Acknowledgment

The Author acknowledges Prof. S. Giller and Prof. P. Kosiński for a careful reading of the manuscript and many helpful suggestion. Special thanks are

also due to Prof. P. Maślanka and Mr. E. Kowalczyk for valuable discussion.

6 Appendix

In this appendix, in order to illustrate the procedures used, we obtain Eqs. (26) for the case 9.

In this case the generating function reads (cf. remark after Eq. (16))

$$\varphi = e^{\mu m} e^{\nu \tau} e^{\kappa v} e^{\rho a}$$

In order to determine coalgebra sector we use the duality relation

$$< \varphi\psi, X > = < \varphi \otimes \psi, \triangle(X) >$$

To this end, we calculate:

$$e^{\mu' m} e^{\nu' \tau} e^{\kappa' v} e^{\rho' a} e^{\mu m} e^{\nu \tau} e^{\kappa v} e^{\rho a} = e^{(\mu' + \mu e^{-i\frac{1}{\chi_1}\varrho'})m} e^{(\nu'+\nu)\tau}$$

$$e^{\left(\kappa e^{-i\frac{1}{\chi_1}\varrho'} + \kappa' \frac{1}{\frac{1}{2} i \frac{1}{\chi_1} \mu e^{-i\frac{1}{\chi_1}\varrho'} v+1}\right) v + (2\frac{\chi_1}{\chi_2}\nu' + 2\frac{\chi_1}{\chi_3}\rho') \ln\left(\frac{1}{2} i \frac{1}{\chi_1} v \mu e^{-i\frac{1}{\chi_1}\rho'} + 1\right)}$$

$$e^{O(v^2,...)} e^{(\rho'+\rho)a}$$

where $O(v^2, ...)$ means function depending on v^2 and higher powers of v.

Taking the first power of v in exponentials we obtain the following result

$$\varphi\psi = e^{(\mu' + \mu e^{-i\frac{1}{\chi_1}\varrho'})m} e^{(\nu'+\nu)\tau}$$

$$e^{\left(\kappa e^{-i\frac{1}{\chi_1}\varrho'} + \kappa' + i\frac{1}{\chi_2}\mu\nu' e^{-i\frac{1}{\chi_1}\varrho'} + i\frac{1}{\chi_3}\mu\rho' e^{-i\frac{1}{\chi_1}\varrho'}\right)v} e^{O(v^2,...)} e^{(\rho'+\rho)a}$$

Therefore we have

$$< \Delta M, e^{\mu' m} e^{\nu' \tau} e^{\kappa' v} e^{\rho' a} \otimes e^{\mu m} e^{\nu \tau} e^{\kappa v} e^{\rho a} > = < M, e^{\mu' m} e^{\nu' \tau} e^{\kappa' v} e^{\rho' a} e^{\mu m} e^{\nu \tau} e^{\kappa v} e^{\rho a} > =$$

$$= -i(\mu' + \mu e^{-i\frac{1}{\chi_1}\varrho'})$$

Using this and the corresponding formulas for H, P and K we arrive at the coalgebra described by Eq.(26)

Now using the duality relation $<\varphi, XY>=<\triangle\varphi, X\otimes Y>$ we can determine the algebra sector. We have:

$$\Delta\varphi = e^{\mu(m'+m-\frac{1}{2}v'^2\tau-av')}e^{\nu(\tau'+\tau)}e^{\kappa(v'+v)}e^{\rho(a'+a+\tau v')}$$

In order to deal with the above expression we first decompose

$$e^{\mu(m'+m-\frac{1}{2}v'^2\tau-av')}$$

To this end let

$$f(\mu) = e^{-\mu m'}e^{\mu(m'+m-\frac{1}{2}v'^2\tau-av')}.$$

Then

$$\Delta\varphi = e^{\mu m'}fe^{\nu(\tau'+\tau)}e^{\kappa(v'+v)}e^{\rho(a'+a+\tau v')}.$$

The function $f(\mu)$ obeys $f(0) = 1$, and

$$\dot{f} = \frac{df}{dx} = \left(m - \frac{1}{2}\frac{\tau v'^2}{(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v')^2} - \frac{av'}{1+\frac{1}{2}i\frac{1}{\chi_1}\mu v'}\right)f$$

The above equation cannot be solved in a standard way due to the fact that the terms appearing on the right hand side do not commute. Therefore we pass to the "interaction picture" by letting

$$f = e^{mX(v',\mu)}h.$$

Then

$$\dot{f} = m\dot{X}e^{mX}h + e^{mX}\dot{h}$$

and we select X in such a way that the terms containing m cancel. It can be checked that X should be of the form

$$X(v',\mu) = \mu(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v').$$

We have now:

$$e^{\mu(m'+m-\frac{1}{2}v'^2\tau-av')}=e^{\mu m'}e^{\mu m(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v')}h$$

Our next aim is calculate h. We do this in the same way by writing out a relevant differential equation

$$\dot{h}=-\left(\frac{av'}{1+\frac{1}{2}i\frac{1}{\chi_1}\mu v'}+\gamma(\tau,v,v',\mu)\right)h$$

and substituting

$$h=e^{\lambda(v',\mu)a}$$

in order to deal with noncommutativity of the terms on right hand side.

Step by step we arrive at following result

$$e^{\mu(m'+m-\frac{1}{2}v'^2\tau-av')}=$$

$$e^{\mu m'}e^{\mu m(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v')}e^{2i\chi_1\ln(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v')a}e^{-\int_0^\mu\gamma(\frac{v}{(1+\frac{1}{2}i\frac{1}{\chi_1}\mu v')^2},v',\tau,\mu)d\mu}$$

where funtion γ is of third degree in v, v'.

Finally, $\Delta\varphi$ can be rewritten as

$$\Delta\varphi=e^{\mu m'}e^{\nu\tau'}e^{\mu m}e^{\frac{1}{2}i\frac{1}{\chi_1}\mu^2 v\prime m}e^{\nu\tau}e^{\kappa v'}e^{O(v'^2,v,\tau)}$$

$$e^{\kappa ve^{i\frac{1}{\chi_1}\mu v'}}e^{-\mu v'a}e^{\varrho a}e^{-i\chi(1-e^{-i\frac{1}{\chi_1}\varrho})v'\tau}e^{\varrho a'}$$

Using the above formula, we may calculate

$$<KM,e^{\mu m}e^{\nu\tau}e^{\kappa v}e^{\rho a}>=<K\otimes M,\Delta(e^{\mu m}e^{\nu\tau}e^{\kappa v}e^{\rho a})>=$$

$$=(-i)(-i)(\frac{1}{2}i\frac{1}{\chi_1}\mu^2)$$

$$<MK,e^{\mu m}e^{\nu\tau}e^{\kappa v}e^{\rho a}>=<M\otimes K,\Delta(e^{\mu m}e^{\nu\tau}e^{\kappa v}e^{\rho a})>=0$$

Therefore:

$$<[K,M],e^{\mu m}e^{\nu\tau}e^{\kappa v}e^{\rho a}>=-\frac{1}{2}i\frac{1}{\chi_1}\mu^2$$

Other commutators are calculated in a similar way.

References

[1] Opanowicz 1998 $J.Phys.A : Math.Gen.$ **31** 2887.

[2] Kowalczyk 1998 q-alg/9804055

[3] Maślanka 1994 $J.Math.Phys.A : Math.Gen.$ **35** 1976.